



AUTHOR CORRECTION

“WHICH PATIENTS SHOULD BE TREATED WITH ANTI-IGE?”. K-M. BEEH. *EUR RESPIR REV* 2007; 16: 104, 85–87.

Unfortunately, the dosage information in figure 1 of the above article was presented incorrectly. The figure should have appeared as seen below.

Baseline IgE·IU mL ⁻¹	Total doses mg per 28 days										
	Body weight kg										
	20–25	>25–30	>30–40	>40–50	>50–60	>60–70	>70–80	>80–90	>90–125	>125–150	
≥ 30–100	75	75	75	150	150	150	150	150	300	300	Q4 weeks
> 100–200	150	150	150	300	300	300	300	300	450	600	Q2 weeks
> 200–300	150	150	225	300	300	450	450	450	600	750	
> 300–400	225	225	300	450	450	450	600	600	Do not administer		
> 400–500	225	300	450	450	600	600	750	750			
> 500–600	300	300	450	600	600	750					
> 600–700	300	450	450	600	750						

FIGURE 1. Omalizumab dosing table (European Union). Ig: immunoglobulin; Q4 weeks: total dose to be administered every 4 weeks; Q2 weeks: half of total dose to be administered every 2 weeks.

DOI: 10.1183/09059180.50010405