

Supplementary Table 2. Radiology features

	Tubbs	Carrington	Yousem	Ryu	Kawabata	Hartman	Akira	Baloira	Individual cases	Total
Chest X-ray present	26 (100%)	40 (100%)	36 (100%)	22 (96%)	30 (97%)				63 (93%)	217
Bilateral interstitial pattern	26 (100%)		32 (89%)	18 (82%)	24 (77%)				53/62 (85%)	153/170 (90%)
Basal predominance					24 (77%)				47/60 (78%)	71/90 (78%)
Normal	0 (0%)	9 (23%)	4 (11%)	2 (9%)					4/63 (6%)	19/187 (10%)
Chest CT present				18 (78%)	26 (84%)	22 (100%)	8 (100%)	11 (92%)	29/63 (45%)	114
Bilateral groundglass				15 (83%)	23 (88%)	22 (100%)	8 (100%)	11 (100%)	26/29 (90%)	105/114 (92%)
Lower lobe predominance						22 (100%)	8 (100%)		24/29 (83%)	54/59 (92%)
Reticulation				3 (17%)		13 (59%)	5 (63%)		10/29 (34%)	31/77 (40%)
Traction bronchiectasis						7 (32%)	1 (13%)	3 (27%)		11/41 (27%)
Cysts					10 (38%)	7 (32%)	8 (100%)	5 (45%)		30/67 (45%)
Honeycombing					5/14 (36%)*		3 (38%)	2 (18%)	2/29 (7%)	12/62 (19%)

Available data from case series and individual cases are presented. Total data are summarized as percentage of available data. * honeycombing developed at follow-up in 5/14 cases who had a follow-up HRCT scan.